

Extraction of the neutron structure function F_2^n from inclusive scattering data on composite nuclei

A.S. Rinat and M. F. Taragin

Weizmann Institute of Science, Department of Particle Physics, Rehovot 76100, Israel

(February 8, 2008)

Abstract

We consider a generalized convolution, linking Structure Functions (SF) F_2^N for nucleons, F_2^A for a physical nucleus and $f^{PN,A}$ for a nucleus, composed of point-nucleons. In order to extract F_2^n we employ data on $F_2^{p,A}$ and the computed $f^{PN,A}$. Only for $Q^2 \approx 3.5 \text{ GeV}^2$ do data permit the extraction of $F_2^A(x, 3.5)$ over a sufficiently wide x -range. Applying Mellin transforms, the above relation between SF turns into an algebraic one, which one solves for the Mellin transform of the unknown F_2^n . We present inversion methods leading to the desired F_2^n , all using a parametrization for $C(x, Q^2) = F_2^n(x, Q^2)/F_2^p(x, Q^2)$. Imposing motivated constraints, the simplest parametrization leaves one free parameter $C(x = 1, Q^2)$. For $Q^2 = 3.5 \text{ GeV}^2$ its average over several targets and different methods is $\langle C(1, 3.5) \rangle = 0.54 \pm 0.03$. We argue that for the investigated Q^2 , $C(x \rightarrow 1, 3.5)$ is determined by the nucleon-elastic (NE) part of SF. A calculation of the latter comes close to the extracted value. Both are close to the $SU(6)$ limit $u_V(x, 3.5) = 2d_V(x, 3.5)$ for parton distribution functions.

It has been recently proposed to extract the ratio $C(x, Q^2) = F_2^n(x, Q^2)/F_2^p(x, Q^2)$ of the

n, p structure functions (SF) from data on ratios $\sigma(^3\text{H})/\sigma(^3\text{He})$ of inclusive cross sections. An electron beam with $E \approx 11\text{--}12$ GeV [1] is expected to provide the required wide, continuous kinematic ranges. The proposal has already elicited discussions, mainly on the stability of the suggested extraction techniques for $C(x, Q^2)$ [2–5].

An experiment, involving the SF of the lightest isobars with minimal I -spin symmetry breaking is presumably best suited to obtain $C(x, Q^2)$. However, even when an upgrading of the beam will be approved, results on the $A=3$ proposal are not expected before 2006. One therefore wonders whether existing data on F_2^A may furnish similar information. Below we explore this possibility and present first results for $C(x, Q^2)$ and $F_2^n(x, Q^2)$ from inclusive scattering data on D, C and Fe.

In the following we limit ourselves to inclusive scattering of unpolarized electrons from randomly oriented targets A . The cross section per nucleon for beam energy E reads

$$\frac{d^2\sigma^{eA}(E; \theta, \nu)/A}{d\Omega d\nu} = \frac{2}{M}\sigma_M(E; \theta, \nu)F_2^A(x, Q^2) \left[\frac{xM^2}{Q^2} + \tan^2(\theta/2) \frac{F_1^A(x, Q^2)}{F_2^A(x, Q^2)} \right], \quad (1)$$

where θ and ν are the scattering angle and the energy loss. $F_{1,2}^A(x, Q^2)$ are nuclear structure functions (SF) per nucleon, expressed in terms of the squared 4-momentum transfer $Q^2 = \mathbf{q}^2 - \nu^2$ and the Bjorken variable $x = Q^2/2M\nu$ (M is the nucleon mass) with range $0 \leq x \leq A$.

Henceforth nuclear and nucleon SF are assumed to be related by some generalized convolution $F^A = f * F^N$, for instance [6,7]

$$F_k^A(x, Q^2) = \int_x^A \frac{dz}{z^{2-k}} \left[\frac{Z}{A} f_p^{PN,A}(z, Q^2) F_k^p(x/z, Q^2) + \frac{N}{A} f_n^{PN,A}(z, Q^2) F_k^n(x/z, Q^2) \right] \quad (2a)$$

$$\approx \int_x^A \frac{dz}{z^{2-k}} f^{PN,A}(z, Q^2) F_k^{(N)}(x/z, Q^2). \quad (2b)$$

In Eq. (2) $f_{p,n}^{PN,A}$ is the SF for a nucleus, composed of point-nucleons, where initially a p or a n absorbs the virtual photon; $f^{PN,A}$ is their average over the number of protons and neutrons in the target $A(Z, N)$. $F_k^{(N)}$ stands for the similarly averaged nucleon (N) SF (We often drop arguments of functions when there is no danger of confusion) ¹.

¹* In Eqs. (2) appear admixtures of nucleon SF [8], which are negligible for the involved Q^2 .

$$\begin{aligned}
f^{PN,A} &= \frac{Z}{A} f_p^{PN,A} + \frac{N}{A} f_n^{PN,A} \\
F_k^{\langle N \rangle} &= \frac{Z}{A} F_k^p + \frac{N}{A} F_k^n
\end{aligned} \tag{3}$$

Eq. (2) describes partons from nucleons in a nucleus, but not those from other sources (virtual bosons) neither does it account for anti-screening. Both limit the use of (2) to $x \gtrsim 0.15$ -0.20 [9]. Eq. (2) is estimated to be valid for $Q^2 \gtrsim Q_c^2 \approx 2 - 2.5 \text{ GeV}^2$ [10,11].

We briefly mention previously suggested extraction methods for F_2^n . Those dealt mostly with D data and use f^{PN} in the Impulse Approximation (IA) [12,13]. It has for instance been emphasized that Eq. (2) is a Fredholm integral equation for the unknown $F_k^{\langle N \rangle}$. Discretization in x and z produces a set of linear equations with a solution, tending to the exact answer for $\Delta z \rightarrow 0$ [14]. The strong variation of the 'kernel' f^{PN} apparently hampers an actual application of the above. We also mention iteration methods to deconvolute nuclear effects for a D target [12,15].

A different deconvolution has been suggested for the proposed precision data, relating F_k^A [1]. For those one may define super-ratios [2,4]

$$\mathcal{R}^{A_1, A_2}(x) \equiv \frac{\rho^{A_1}(x)}{\rho^{A_2}(x)} \tag{4a}$$

$$\rho^A(x) = \frac{[F_2^A(x)/F_2^p(x)]}{a[F_2^n(x)/F_2^p(x)] + b} , \tag{4b}$$

with, in principle, arbitrary a and b . For given F_2^p those ratios contain $C(x)$ explicitly, and are from Eq. (2) seen to depend implicitly on $C(x/z)$. Iteration determines $C(x)$.

Finally we recall the expansion of $F^{\langle N \rangle}(x/z)$ in Eq. (2) around the maximum $z \approx 1$ of $f(z)$ [16]

$$F_2^A(x) = F_2^{\langle N \rangle}(x) + \mathcal{M}_1(x)[x F_2^{\langle N \rangle(1)}(x)] + \mathcal{M}_2(x)[x F_2^{\langle N \rangle(1)}(x) + \frac{x^2}{2} F_2^{\langle N \rangle(2)}(x)] + \dots , \tag{5}$$

with $\mathcal{M}_n(Q^2) = \int_0^A dz f^{PN,A}(z, Q^2)(1-z)^n$. The first A -independent term $F_2^{\langle N \rangle}(x)$ adequately represents the series for $x \lesssim 0.4$. Insufficient knowledge of $F_2^{\langle N \rangle(l)}(x)$, the l -th derivative of $F_2^{\langle N \rangle}(x)$, prevents the use of the expansion (5) beyond that range. Although not the region of prime interest, the low- x result will guide the extraction of F_2^n .

We first recall calculations of F_k^A from Eq. (2) with $F_k^{p,n}$ as input. For nucleons in vacuum there are abundant data on F_2^p [17] and less accurate ones for F_1^p [18]. Inclusive data do not reach the elastic point $x = 1$, but the used parametrizations do. With no direct information on F_k^n , one frequently uses the 'primitive' assumption

$$F_k^{n,pr} \equiv 2F_k^D - F_k^p, \quad (6)$$

which is the first term in Eq. (5) for the D and thus holds only for small x . The question of interest is how modifications of Eq. (6) for $x \gtrsim 0.4$ influence $F_k^A(x, Q^2)$, Eq. (2).

The SF $f^{PN,A}$ is a many-body property and requires a model for a calculation, such as the perturbative IA, which to lowest order contains the single-hole spectral function [19,20]. There nucleons appear off their mass shell and a prescription is needed to handle those. We prefer a non-perturbative version, based on a generalized Gersch-Rodriguez-Smith (GRS) theory, where the SF of nucleons F^N are reasoned to be on their mass shell. That method moreover allows a computation of $f^{PN,A}$ beyond its lowest order [7,11,21–23].

The above program has initially been realized for $A \geq 12$. It has been shown that $f^{PN,A}$ is nearly independent of A , and using Eq. (2) one proves the same for F_k^A . Data indeed show that for $x \lesssim 0.8$, the ratios $\mu^{A,A'} = [d^2\sigma^{eA}/A]/[d^2\sigma^{eA'}/A'] \approx F_2^A/F_2^{A'}$, $A, A' \geq 12$ differ from 1 by less than 2-3% [21,23,24].

The above does not hold for the lightest nuclei, in particular not for $A' \rightarrow D$, when

$$\mu^A(x, Q^2) = \mu^{A,D}(x, Q^2) = \frac{d^2\sigma^{eA}(x, Q^2)/A}{d^2\sigma^{eD}(x, Q^2)/2} \approx \frac{F_2^A(x, Q^2)}{F_2^D(x, Q^2)} \quad (7)$$

are the EMC ratios, with their characteristic deviations from 1 in the range $x \lesssim 0.9$ [25].

For $A \leq 4$ and given NN interaction, one may presently compute with great precision nuclear ground states [26], as well as $f^{PN,A}$ in Eq. (2). Using those, GRS calculations of inclusive cross sections have recently been completed for D [27] and ${}^4\text{He}$ [28].

We now address the inverse problem of trying to extract F_2^n from data on $F_2^{p,A}$ and the above-mentioned computed f . First one needs to obtain nuclear SF and we shall use JLab data for D [29,30] and for C, Fe [31], supplemented by some older NE3 data of more restricted kinematics [32]. We mention two approaches:

i) *R*-ratios: Ideally one performs a Rosenbluth extraction of F_k^A from cross sections (1) for fixed x, Q^2 and varying θ or E , which provides

$$R^A(x, Q^2) = \left(1 + \frac{4M^2x^2}{Q^2}\right) \left(\frac{F_2^A(x, Q^2)}{2xF_1^A(x, Q^2)}\right) - 1 \quad (8)$$

Unfortunately, the set of (x, Q^2) points in the JLab and NE3 data does not enable the above Rosenbluth extractions. Instead one frequently invokes an empirical expression such as $R^A(x, Q^2) \approx 0.32/Q^2$. It assumes A , as well as x -independence and prescribes the dependence on Q^2 [33]. The above empirical R actually contradicts theoretical predictions [11]. We shall return to this point below.

ii) *Theoretical information*: Using the primitive $F^{n,pr}$, computed cross sections (1) as function of θ, ν generally agree with data in the deep-inelastic regions, $x \lesssim 1$, ($Q^2 \gtrsim Q_c^2$). There we select data, for which the relative deviation $\alpha(= \alpha(x, Q^2)) = (F_2^{A,th} - F_2^{A,exp})/F_2^{A,exp}$ satisfies $|1 - \alpha| \lesssim 0.2$ and which, for given θ appear to change smoothly with x . We suggest to attribute the above deviations in equal measure to both structure functions F_k^A in (1), which enables the definition of quasi-data

$$F_k^{A,qd}(x, Q^2) \equiv \alpha(x, Q^2) F_k^{A,th}(x, Q^2) \quad (9)$$

By construction, the $F_k^{A,qd}$ produce an exact fit to cross sections.

Use of Eq. (2) for the purpose of inversion requires input for fixed Q^2 and running x , whereas cross section data are for essentially one beam energy E and cover separated x bins. Q^2 varies mildly within those x -bins, but not when going from one bin to a neighbouring one. It is thus necessary to make cuts for fixed Q^2 , containing a maximal number of x -points with $x \lesssim 1.0 - 1.1$. After careful interpolation between quasi-data, the above appears only fulfilled for Q^2 between 3-4 GeV² and then only for $0.5 \lesssim x \lesssim 1.1$. It is clearly impossible to reliably extrapolate the manipulated quasi-data into the crucial range $x \lesssim 0.5$.

At this point we invoke the empirical observation that for $x_0 \approx 0.16 - 0.18$, $F_2^p(x_0, Q^2)$ and $F_2^D(x_0, Q^2)$ are practically constant as function of Q^2 [17]. We exploit this by considering Eq. (2) for small $x \leq x_0$. Upon substitution of the small- x part of Eq. (5) into (2) and use of $\int_0^A dz f^{PN,A}(z) = 1$, one easily shows that for *any* target

$$F_2^{A,nucl}(0.18) \approx F_2^N(0.18) \approx 0.34 \quad (10a)$$

$$F_2^{A,nucl}(0) = F_2^{\langle N \rangle}(0) = F_2^N(0) \neq 0 \quad (10b)$$

The latter extension to $x = 0$ is in agreement with Eq. (5) and will be needed below ².

We recall that $f^{PN,A}$ in Eq. (2) relates only to nucleons in the target and as a result, Eqs. (10a) hold only for the nucleonic component $F_2^{A,nucl}$. That component may now be *interpolated* for $x \lesssim 0.55$, completing knowledge of $F_2^{A,nucl}(x)$ over the entire relevant range $x \lesssim 1.1$.

After the critical remarks on the empirical form of the R -ratio it comes as a surprise that in the range $0.5 \lesssim x \lesssim 1$, $F_2^{Fe,qd}(x, 3.5)$ and the empirically extracted $F_2^{Fe(R)}(x, 3.5)$ agree to within $\pm 5\%$, and even for $1 \lesssim x \lesssim 1.5$ to within $\pm 12\%$! It vindicates Arrington's claim that even a 100% uncertainty in the phenomenological R incurs only a 5% uncertainty in $F_2^{A(R)}$ [30].

We proceed as follows. For Fe, for which the number of available deep-inelastic data points is largest, we use $F_2^{Fe,qd}$ from method ii). With insufficient C, D data for application of that method, we exploit Eq. (7) and find

$$\begin{aligned} F_2^{C,qd} &\equiv \frac{F_2^{C,R}}{F_2^{Fe,R}} F_2^{Fe,qd} \\ F_2^{D,qd} &= [\mu^C]^{-1} F_2^{C,qd} \end{aligned} \quad (11)$$

EMC data for μ^C are in the desired range $0.3 \lesssim x \lesssim 0.80$ where they require some smoothing. For $x \lesssim 0.30$ we interpolated the purely nucleonic component as in ii) and added for $0.80 < x < 0.88$ averages of Be and Al data [25].

We now introduce Mellin transforms (MT) and their inverses, which for real $g(x)$ are defined as

$$\tilde{g}(u) = \int_0^\infty dx x^{u-1} g(x) \quad (12a)$$

$$g(x) = \frac{1}{\pi} \int_0^\infty dt \operatorname{Re} \left[\frac{g(a+it)}{x^{a+it}} \right], \quad (12b)$$

^{2*} The above explains why for all A and Q^2 , EMC ratios $\mu^A(x, Q^2)$ cross 1 for $x \approx 0.18$ [25].

The constant a is chosen such, that $g(u) = g(a + it)$ is free of singularities in the complex u plane to the right of the imaginary u axis shifted by a .

Application of Eq. (12a) to Eq. (2), turns the generalized convolution into a linear relation, which can be solved for $\tilde{F}_2^n(u, Q^2)$. For fixed Q^2

$$\tilde{F}_2^A(u) = \tilde{f}^{PN,A}(u+1)\tilde{F}_2^{(N)}(u) + \mathcal{G}^{(N(NE))} \quad (13a)$$

$$\tilde{\mathcal{F}}_2^{n(A)}(u) = \frac{A}{N} \frac{\tilde{F}_2^A(u)}{\tilde{f}^{PN,A}(u+1)} - \frac{Z}{N} \tilde{F}_2^p(u) - \left[\frac{Z}{N} \mathcal{G}^p + \mathcal{G}^n \right], \quad (13b)$$

with $\mathcal{G}^{N,NE}$, combinations of the standard nucleon static formfactors, making up the NE parts of $F_2^{N,NE}$ (see for instance Ref. [11]). Above we denote by $\tilde{\mathcal{F}}_2^{n(A)}$ the $A(N, Z)$ -dependent right-hand side of Eq. (13b). For exact input F_2^A , it should coincide with $\tilde{F}_2^n(u)$, the MT of $F_2^n(x)$.

Next we consider $u = 0$ in Eq. (13b). Since the normalization of $f^{PN,A}$ implies $\tilde{f}^{PN,A}(1) = 1$, one finds

$$\tilde{\mathcal{F}}_2^{n(A)}(u=0) = \frac{A}{N} \tilde{F}_2^{A,nuc}(u=0) - \frac{Z}{N} \tilde{F}_2^p(u=0) - \left[\frac{Z}{N} \mathcal{G}^p + \mathcal{G}^n \right] \quad (14)$$

Consequently, Eq. (12a), when used in (10a) implies that none of the MT of SF are defined for $u = 0$. This has numerical consequences also in the immediate neighbourhood of $u = 0$.

Unfortunately, the direct inversion Eq. (12b) of $\tilde{\mathcal{F}}(u)$ runs into serious numerical problems. We therefore take recourse to indirect methods, all featuring a parametrization of the ratio C in

$$F_2^n(x, Q^2) = F_2^n(x, Q^2; d_k) = C(x, Q^2; d_k) F_2^p(x, Q^2) \quad (15)$$

$$C(x, Q^2; d_k) = \sum_{k \geq 0} d_k(Q^2) (1-x)^k$$

Eq. (10a) implies a first constraint $C(0) (= F_2^n(0)/F_2^p(0)) = \sum_{k \geq 0} d_k(Q^2) = 1$. Next, one may exploit $F_2^n(x) = F_2^{n,pr}(x)$, Eq. (5), for small $x \lesssim 0.35$. We use only $F_2^n(0.2, 3.5) = 0.75$ which for $k_{max} = 2$ leaves one free parameter $d_0 = C(1)$.

It is convenient to re-parametrize $F_2^p(x, 3.5)$ as follows

$$F_2^p(x, Q^2) = x^{-a^2} \sum_{m \geq 1} c_m (1-x)^m; x \geq 0.02 \quad (16a)$$

$$= 0.42 \quad ; x \leq 0.02 \quad (16b)$$

A small $a^2 \ll 1$ in Eq. (16a) produces the observed rise of $F_2^p(x)$ for small x , while the cut-off in Eq. (16b) avoids the singularity in (16a). Eq. (10a) again implies the same for $F_2^{A,n}(x)$. In the region $0.02 \lesssim x \lesssim 0.9$, F_2^p , Eq. (16a), practically coincides with the parametrization in Ref. [17], both agreeing with data averaged over resonances.

We now explore some extraction methods, comparing relevant computed or extracted expressions A , related to F_2^n and parametrized forms B :

I) The extracted $\tilde{\mathcal{F}}_2^{n(A)}(u)$, Eq. (13b) and the MT of F_2^n , Eq. (15).

II) $F_2^{A,qd}(x)$, Eq. (9), with the nuclear SF $F_2^A(x)$, computed from (2), now using Eq. (15).

The parameters d_k , are determined by minimization of variances $w(d_0) = \sum_i |A_i - B_i|^2$ (or of relative variances). Table I summarizes our results. The spread of the individual entries $C(1)$ from I) reflects variations in the smallest and largest u , retained in the above sum (see remark after Eq. (14)). An empty entry indicates the absence of a minimum in the studied intervals. We note that in those cases the slope of the variance is very small and that the range of $C(1)$ is actually compatible with values from real minima. The results for the two methods and for the three targets produces a well-determined average $\langle C(1, 3.5) \rangle = 0.54 \pm 0.03$.

In Fig. 1 we show the full $C(x, 3.5)$, as well as $F_2^{p,n}(x, 3.5)$. On the right abscissa are marked the values $2/3, 3/7, 1/4$ for $C(1)$, corresponding to exact SU(6) symmetry, dominance of $S = 0$ over $S = 1$ di-quark coupling and the same for the z -component S_z for $x \rightarrow 1$ (see for instance Ref. [15])³.

We now argue that for our purposes the regarded $Q^2 \approx 3.5 \text{ GeV}^2$ is not 'large'. First

^{3*} The extracted F_2^n allows an evaluation of the Gottfried sumrule $S_G(3.5) = \int_0^1 (dx/x) [F_2^p(x, Q^2) - F_2^n(x, Q^2)] = 0.251$. This is close to the recent value $S_G(4.0) = 0.256 \pm 0.026$

we note that $F_2^{p,n}(x, Q^2)$ has a first inelastic threshold $N + m$, m being the pion mass, or $x_{thr}(Q^2) = [1 + 2Mm/Q^2]^{-1}$, i.e. a finite x -distance from the elastic point $x = 1$. That interval shrinks with increasing Q^2 , but for the case at hand $x_{thr}(3.5) \approx 0.93$, which is marked by a vertical line in Fig. 1. Data show that $F_2^p(x, 3.5)$ is negligibly small beyond $x \approx 0.9$ [12,29,34].

The above implies that $C(x \rightarrow 1, 3.5)$ can be ascribed to the NE part $F^{N,NE}$, i.e. to static N form factors (cf. Eq. (13b)). Disregarding the electric form factor G_E^n of the n , one finds (μ_N are the magnetic moments of the nucleons)

$$\lim_{x \rightarrow 1} C(x, Q^2) = \left[\frac{\mu_n \alpha_n(Q^2)}{\mu_p \alpha_p(Q^2)} \right] \left[1 + \frac{4M^2}{Q^2} \left(\frac{\gamma(Q^2)}{\mu_p} \right)^2 \right]^{-1} \quad (17)$$

with

$$\gamma(Q^2) = \frac{\mu_p G_E^p(Q^2)}{G_M^p(Q^2)}; \quad \frac{\alpha_n(Q^2)}{\alpha_p(Q^2)} = \frac{G_M^n(Q^2)/\mu_n}{G_M^p(Q^2)/\mu_p} \quad (18)$$

Recent data show that $\gamma, \alpha_p, \alpha_n$ deviate from 1 and equal for $Q^2 = 3.5, 5.0 \text{ GeV}^2$: $\gamma=0.552$, 0.349 and $\alpha_n/\alpha_p \approx 1.2, 1.1$ [35,36]. Eqs. (17), (18) then yield

$$C(x = 1, 3.5) \approx 0.61; \quad C(x = 1, 5.0) \approx 0.56 \quad (19)$$

The large Q^2 limit essentially depends on the same for α_n/α_p . For a value 1, $C(x \rightarrow 1, Q^2 \rightarrow \infty) = 0.469$, which shows that in the above sense $C(1, 3.5)$ is still far from a scaling limit.

The above value 0.61 is reasonably close to the extracted one and has also been entered on the abscissa in Fig.1. Either one definitely exceeds previously cited values $C(1) \approx 0.42$ from D data (cf. Ref. [15]). The above seems to indicate that SU(6) symmetry $u_v(x, 3.5) = 2d_v(x, 3.5)$ for the up and down quark parton distribution functions is only mildly broken. It is moreover in agreement with globally extracted distribution functions [38,39].

[37], because of nearly identical contributions from the dominant small x region. We note in passing that a finite outcome requires $C(0, Q^2) = 1$. For CTEQ parametrizations from global parton-distribution functions $C(0, Q^2) = (1 + \delta C(0))$ [38], leading to a diverging S_G .

The above reasoning rests on forms for the nucleon SF, averageKd over resonances. Retaining those in detail, the sharp first inelastic threshold will become blurred, but it seems reasonable that $C(1, Q^2)$ for relatively low Q^2 remains determined by static elastic form factors. That point is presently under study. A similar large- Q^2 result has been found from quark-hadron duality considerations, pushed to the extreme for the part $F_2^{N(NE)}$ [40].

Summarizing, we have described methods to extract the neutron Structure Function from existing inclusive scattering data on various targets. The experimental material from several targets provided consistent values for $C(1, 3.5)$. It is very desirable to plan the forthcoming inclusive scattering experiments on D and ^4He with the available 6 GeV beam [41], such that there be coverage of continuous $x \lesssim 1$ -range for at least one Q^2 . Confrontation of those data with accurate calculations, possible for those nuclei [27,28] will sharpen the present outcome for $F_2^n(x, Q^2), C(x, Q^2)$.

The authors thank G. Petratos for giving detailed information on the proposed $A=3$ experiment and J. Arrington for putting at our disposal D and Coulomb-corrected NE3 data for C and Fe. ASR profited from discussions with G. Salmè, E. Pace, several experimentalists at JLab and in particular with W. Melnitchuk .

REFERENCES

- [1] G.G. Petratos, Draft version PAC18, Jefferson Lab, July 2000.
- [2] I.R. Afnan, F. Bissey, J. Gomez, A.T. Katramatou, W. Menitchouk, G.G. Petratos and A.W. Thomas, Phys. Lett. B 493 (2000) 36; F. Bissey, A.W. Thomas and I.R. Afnan, Phys. Rev. C 64 (2001) 024004.
- [3] K. Saito, C. Boros, K. Tsushima, F. Bissey, I.R. Afnan and A.W. Thomas, Phys. Lett. B 493 (2000) 288.
- [4] E. Pace, G. Salmè, S. Scopetta and A. Kievsky, Phys. Rev. C64 (2000)05523; Nucl. Phys. A 689 (2001) 453.
- [5] M.M. Sargassian, S. Simula and M.I. Strickman, nucl-th/0105052
- [6] S.V. Akulinichev, S.A. Kulagin and V.M. Vagrado, Phys. Lett. 158 B (1985) 485; G.V. Dunne, Nucl. Phys. A455 (1986) 701.
- [7] S.A. Gurvitz and A.S. Rinat, TR-PR-93-77/ WIS-93/97/Oct-PH; Progress in Nuclear and Particle Physics, Vol. 34 (1995) 245.
- [8] G.B. West, Ann. of Phys. (NY) 74, (1972) 464; W.B. Atwood and G.B. West, Phys. Rev. D 7, (1973) 773.
- [9] C.H. Llewelyn Smith, Phys. Lett B 128 (1983) 107; M. Ericson and A.W. Thomas, *ibid* 112.
- [10] A.S. Rinat, Proceedings of Meeting 'Prospects on Hadron and Nuclear Physics', Trieste, IT (1999) World Scientific.
- [11] A.S. Rinat and M.F. Taragin, Phys. Rev. C 62 (2000) 034602.
- [12] A. Bodek *et al*, Phys. Rev. D 20 (1979) 1471.
- [13] S. Liuti and Franz Gross, Report TH-95-06

- [14] A.Yu. Umnikov, F.C. Khanna and L.P. Kaptari, Z. F. Physik A 348 (1994) 211.
- [15] W. Melnitchouk and A.W. Thomas, Phys. Lett. B 377 (1996) 11.
- [16] S.V. Akulinitchev, S.A. Kulagin and G.M. Vagradov, Phys. Lett. B158 (1985) 485.
- [17] P. Amadrauz *et al*, Phys. Lett B295 (1992) 159; M. Arneodo *et al*, *ibid* B364 (1995) 107.
- [18] A. Bodek and J. Ritchie, Phys. Rev. D 23 (1981) 1070.
- [19] See for instance: C. Ciofi degli Atti, E.Pace and G. Salmè, Phys. Rev. C 43 (1991) 11275; C. Ciofi degli Atti, D.B. Day and S. Liuti, *ibid* C 46 (1994) 1045.
- [20] O. Benhar, A. Fabrocini, S. Fantoni, G.A. Miller, V.R. Pandharipande and I. Sick, Phys. Rev. C44 (1991) 2328; Phys. Lett. B359 (1995) 8.
- [21] A.S. Rinat and M.F. Taragin, Nucl. Phys. A598 (1996); 349 *ibid* A620, 412 (1997); Erratum *ibid* A623 (1997) 773.
- [22] S.A. Gurvitz and A.S. Rinat, nucl-th/0106032; Phys. Rev. C 65, to be published.
- [23] A.S. Rinat and M.F. Taragin, Phys Rev. C 60 (1999) 044601.
- [24] J.Arrington *et al*, Phys. Rev. C 53 (1996) 224
- [25] J. Gomez *et al*, Phys. Rev. D 49 (1994) 4348.
- [26] H. Kamada *et al*, Phys. Rev. C 64 (2001) 044001.
- [27] A.S. Rinat and M.F. Taragin, Phys. Rev. C 65 (2002) 041610.
- [28] M. Viviani, A. Kievsky and A.S. Rinat, nucl-th/0111048, submitted to Phys. Rev. C
- [29] I. Niculescu *et al*, Phys. Rev. Lett. 85 (2000) 1182.
- [30] J. Arrington, private communication.
- [31] J. Arrington *et al*, Phys. Rev. Lett. 82 (1999) 2056; CalTech PhD thesis 1998.

- [32] D.B. Day *et al*, Phys. Rev. C 40 (1993) 1849.
- [33] S. Dasu *et al* Phys. Rev. Lett. 61 (1988), 1061; Phys. Rev. D 49 (1994) 5641; P.E. Bosted *et al*, Phys. Rev. C 46 (1992) 2505; B.W. Fillipone *et al*, Phys. Rev. C 45 (1992) 1582.
- [34] I. Niculescu *et al*, Phys. Rev. Lett. 85 (2000) 1186.
- [35] M. Jones *et al*, Phys. Rev. Lett. 84 (2000) 1398; Third Workshop on 'Perspective in Hadronic Physics' Trieste 2001, IT; to be published.
- [36] A.F Sill *et al*, Phys. Rev. D 48 (1993) 29.
- [37] M. Arneodo *et al*, Phys. Rev. D 50 (1994) R1.
- [38] H.L. Lai *et al*, Phys. Rev. D 51 (1995) 4763; *ibid* D 55 (1997) 1280; Eur. Phys. J. C 12 (2000) 375.
- [39] V. Barone, C. Pascaud and F. Zomer, Eur. Phys. J. C 12 (2000) 243,
- [40] W. Melnitchouk, Phys. Rev. Lett. 86 (2001) 35.
- [41] J. Arrington *et al*, Spokesman proposal JLab, May 2000.

Figures.

Fig. 1. The ratio $C(x, 3.5) = F_2^n(x, 3.5)/F_2^p(x, 3.5)$ for $Q = 3.5 \text{ GeV}^2$ from data on D, C, Fe. The drawn line corresponds to $C(1) = 0.54$ and the band represents the spread in the result, from averaging over different targets and methods. The vertical line for $x = 0.93$ marks the pion threshold $x_{thr}(3.5)$. The numbers on the right abscissa are quark model and QCD predictions for $C(1)$, while 0.61 is the NE limit (18). Also entered are F_2^p and F_2^n , corresponding to C . The band for the latter is hardly noticeable in F_2^n .

TABLES

TABLE I. Values of $C(1,3.5)$ from minimizing variances $w(d_0)$. Results are from data on D,C,Fe, using extraction methods I),II). The spread in results for I correspond to varying u -intervals in the variances. No entry corresponds to cases for which there is no minimum within the above intervals.

	D	C	Fe
I	0.55 ± 0.05	-	0.55 ± 0.03
II	0.55	0.50	-

